

Acupuncture – self-appraisal and the reward system

Thomas Lundeberg, Iréne Lund, Jan Näslund

Abstract

Acupuncture is an ancient therapy with a variety of different explanatory models. A cascade of physiological effects has been reported, both in the peripheral and the central nervous system, following the insertion of a needle or light tapping of the skin. Clinical trials testing the specific claims of acupuncture have generally tried to focus on testing the efficacy of applying specific techniques and/or specified points. However, different conditions may respond differently to different modes of stimulation.

Recently, it was demonstrated that both superficial and deep needling (with *de qi/Hibiki*) resulted in amelioration of patellofemoral pain and unpleasantness. The pleasurable aspect of the acupuncture experience has largely been ignored as it has been considered secondary to its pain alleviating effects. This aspect of acupuncture treatment is likely to be related to activation of self-appraisal and the reward system.

When a patient seeks a therapist there are expectations of a specific effect. These expectations are partly based on self-relevant phenomena and self-referential introspection and constitute the preference. Also, when asked about the effect of the treatment, processes that orientate pre-attentive anticipatory or mnemonic information and processes that mediate self-reflection and recollection are integrated together with sensory detection to enable a decision about the patient's perception of the effect of acupuncture treatment. These 'self-appraisal' processes are dependent on two integrated networks: a ventral medial prefrontal cortex–paralimbic–limbic 'affective' pathway and a dorsal medial prefrontal cortex–cortical–hippocampal 'cognitive' pathway.

The limbic structures are implicated in the reward system and play a key role in most diseases and illness responses including chronic pain and depression, regulating mood and neuromodulatory responses (eg sensory, autonomic, and endocrine). The pleasurable and neuromodulatory aspects of acupuncture as well as 'placebo needling' may partly be explained by the activation or deactivation of limbic structures including the hippocampus, amygdala, and their connections with the hypothalamus.

In patients with patellofemoral pain, the effects of superficial and deep needling remained for six months. These long term pain-alleviating effects have been attributed to activation of pain inhibiting systems in cortical and subcortical pathways. When considering long term effects the cortical–cerebellar system needs to be taken into account. The cortical–cerebellar system is probably central to the development of neural models that learn and eventually stimulate routinely executed (eg motor skills) and long term (eg pain alleviation) cognitive processes. These higher order cognitive processes are initially mediated in prefrontal cortical loci but later shift control iteratively to internal cerebellar representations of these processes. Possibly part of the long term healing effects of acupuncture may be attributed to changes in the cerebellar system thereby sparing processing load in cortical and subcortical areas.

As cortical and subcortical structures are activated and/or de-activated following stimulation of receptors in the skin, disregarding site, 'placebo or sham needling' does not exist and conclusions drawn on the basis that it is an inert control are invalid.

'Self' may be seen as a shifting illusion, ceaselessly constructed and deconstructed, and the effect of acupuncture may reflect its status (as well as that of the therapist).

Keywords

Acupuncture, cerebellum, depression, limbic structures, medial prefrontal cortex, pain, placebo control, reward system, self-appraisal.

Thomas Lundeberg
senior consultant
Rehabilitation Medicine
University Clinic
Danderyds Hospital
Stockholm, Sweden

Iréne Lund
registered
physiotherapist

Jan Näslund
registered
physiotherapist
Department of
Physiology and
Pharmacology
Karolinska Institutet
Stockholm, Sweden

Correspondence:
Thomas Lundeberg

thomas.lundeberg@ds.se

Introduction

Acupuncture is an ancient therapy with different explanatory models.¹ A number of physiological effects have been reported, both in the peripheral and the central nervous systems, following the insertion and stimulation of a needle. Trials testing the specific claims of acupuncture have generally tried to focus on testing the efficacy of varying stimulation techniques and/or specified points. However, insertion of needles into the body can stimulate specific effects though these are not dependent on the locations of stimulation.² Therefore, factors that are simultaneously site specific and unrelated to the site of needling might play a role in the outcome of acupuncture therapy for pain. It is also important to note that different pain conditions might respond differently to different modes of stimulation.³

Patellofemoral pain syndrome (PFPS) is one of the most common musculoskeletal disorders, especially in young adults.⁴ It has been suggested that patients suffering from PFPS are more prone to develop knee osteoarthritis. However, there is no consensus on the definition of PFPS, its classification, assessment, diagnosis, or management. Symptoms and clinical findings in subgroups of individuals with PFPS, classified on the basis of the findings in radiological examinations and compared with subjects with healthy knees, have recently been reported.⁴ An orthopaedic surgeon and a physical therapist consecutively examined 75 patients clinically diagnosed as having PFPS. Radiography revealed pathology in 15 patients, and scintigraphic examination revealed focal uptake of technetium in two patients indicating local pathology. Diffusely increased uptake (sign of sympathetic hyperactivity) was present in 29 patients. In the remaining 29 patients, radiographic and scintigraphic examinations were normal. Subjects with healthy knees differed significantly from the three patient groups in all clinical tests measuring pain in response to provocation: ie compression test, medial and lateral tenderness and passive gliding of the patella. Differences in clinical tests between the patient groups were not significant. The main finding and conclusion of the study was that, in patients with PFPS, it was not possible to detect peripheral pathology with the tests used and that the pain might be due to sympathetic hyperactivity.

A randomised controlled study was conducted to evaluate the effect of acupuncture treatment of patients with PFPS.⁵ Fifty eight patients, clinically and radiologically examined, were randomly assigned to either deep acupuncture treatment with low frequency electrical stimulation or minimal superficial acupuncture with attached electrodes but without electrical stimulation. The patients were treated twice weekly for a total of 15 treatments. The main outcome measurements were one leg vertical jump, functional score, daily visual analogue scale (VAS) recordings of perceived pain intensity, and skin temperature. Fifty-seven patients completed the study. After the treatment period, pain intensity was rated significantly decreased within both groups but with no significant difference between the groups. The decline in pain recordings remained significant even after three and six months in both groups. Even though the pain was rated as decreased after sensory stimulation, the ability to jump on one leg, the functional score, and the skin temperature remained unchanged. This study showed that patients with PFPS appear to benefit from both electroacupuncture treatment and subcutaneous needling. The pain relieving effect of acupuncture remained for six months. The effects are likely to be attributable to activation of pain inhibitory mechanisms or deactivation of pain processes in the central nervous system. Also, it is generally observed following acupuncture that many patients experience reduction of unpleasantness, restoration of wellbeing (health) and their sense of self.

Recently, Hui using accumulative neuroimaging results, thereby achieving larger sample sizes than previously obtained from a single trial, reported that superficial tactile stimulation results in changes in a subset of deep brain structures, although of weaker magnitude and more limited in extent than real acupuncture (matched for acupuncture points and subjects).⁶ The difference between real acupuncture and tactile stimulation varied with the acupuncture points, which may be attributed to differences in innervation of the stimulated tissue. This is supported by Wu and collaborators who found that superficial needling produces many of the sensations elicited with 'real' acupuncture needling, but at a lower frequency and with a different pattern of distribution.⁷ It has been reported that superficial needling results in the activation of A-beta and A-delta nerve fibres

inducing analgesia.⁸ One reason why previous studies on acupuncture neuroimaging have not reported on the effects of superficial acupuncture is possibly related to the fact that these studies have been underpowered to reveal the more subtle actions of superficial needling. Moreover, many of the limbic structures located at the base of the brain such as the amygdala are vulnerable to susceptibility artefacts and suffer signal loss in studies that use thicker sections in the axial orientation (Hui KKS, personal communication).

Acupuncture and self-appraisal

In many patients, reduction of unpleasantness, restoration of wellbeing (health) and the patient's sense of self may be of greater importance than the actual reduction in pain intensity. Neuroimaging studies have shown that there are dedicated brain systems central to appraising the self-relevant content of one's environment and one's conscious mental events.⁹ These systems are adjusted to the detection of environmental phenomena (exteroceptive and interoceptive information: sensory, somatic, autonomic) that convey significance to a patient. Also, the systems are able to install self-relevant information (introspective information: thoughts, memories) from past experience to create abstract associations with stimuli, or, to initiate such information independent of external stimuli altogether (for example expectancy). In this way, one may consider salience simply as propagation of sensory information (like acute pain) sufficient for attention, whereas wellbeing and self-relevance are central processes generated by at least two integrative sub-systems: one that orientates pre-attentive information to self-relevant phenomena and one that engages pensive processes like self-reflection.⁹⁻¹⁵

The task domains that orient preventative or mnemonic information (pre-attentive biasing information) to salient or openly self-relevant phenomena fall into the broad cognitive-affective domains of reward, fear, pain, and affection. The neural substrata include the ventral medial prefrontal cortex (vMPFC), anterior cingulate (ACC), nucleus accumbens (Nacc, also denoted ventral striatum), amygdala (Amg), and insula. The vMPFC is central for detection of self-relevant information (prepotent, biasing or self-relevant) whereas the primary function of Amg and Nacc (limbic structures) is discrimination

of a neuromodulatory response (sensory, autonomic, and endocrine) to specific biasing features of the tasks to facilitate heightened watchfulness of immediately or potentially self-relevant environmental cues.¹⁶⁻⁵⁰

When patients are asked how an acupuncture treatment (stimulus) makes them feel (self-relevant tasks) there is a shift to one's self as the referent which results in activity in distinct neural structures including the ventral and dorsal medial prefrontal cortex (dMPFC), dorsorostral (rACC), and posterior cingulate (PCC). These findings have implications for acupuncture research, suggesting that so called 'placebo acupuncture' is not inert as the patients are continuously asked how the stimulation makes them feel. Also, treatments (stimuli) that convey general information for survival or wellbeing are related to activation in the vMPFC and ACC, Amg, Nacc and insula. Taken together, detection of self-relevance from exteroceptive or interoceptive inputs (needle manipulation and being asked about the treatment effect) triggers a cascade of subcortical processing that orientates the subject or patient on to an increased response potential. Re-integration of this information at vACC and MPFC loci prevent or diminish redirection of attention based on prior experience through integration with contextually appropriate memory traces, of one's past experiences. This would suggest that when given a treatment (drug or acupuncture), the specific meaning of the treatment results in activation of associated self-relevant processes ie an intervention for pain alleviation results in pain alleviation and a drug for insomnia results in induction of sleep.⁵¹⁻⁷⁷

The neural substratum that engages introspective processes includes the dorsal-ventral MPFC, dorsorostral ACC and PCC.⁹ Specifically, this response pattern has been observed during self-appraisal of one's own personality traits, as well as one's opinions, personal morals, attitudes, and during appraisal of one's own preference (eg food, colour). PCC is involved in retrieval of personally significant memories (episodic information) and the dorsorostral ACC function in executive control processes such as selective attention. In contrast to the attention allocating mechanism of the dorsorostral ACC, the dorsal anterior MPFC is involved in evaluative processing of self-relevant mental content. This distinction between anterior MPFC and dorsorostral

ACC function is in line with conclusions drawn from a recent meta-analysis on social cognition, wherein the prior structure was implicated in self-knowledge, and evaluative processes in selective attention.

When a patient seeks a therapist there are expectations of a specific effect. These expectations are partly based on self-relevant phenomena and self-referential introspection and constitute the preference. Also, when asked about the effect of the treatment, processes that orientate pre-attentive anticipatory or mnemonic information and processes that mediate self-reflection and recollection are integrated together with sensory detection to enable a decision about the patient's perception of the effect of acupuncture treatment. These 'self-appraisal' processes are dependent on two integrated networks: a ventral medial prefrontal cortex–paralimbic–limbic 'affective' pathway and a dorsal medial prefrontal cortex–cortical–hippocampal 'cognitive' pathway.^{9,68}

Pariante and collaborators explored the cerebral consequences of needling and expectation with 'real' acupuncture, 'placebo control' acupuncture (Streitberger needle – non-penetrating, blunt telescopic needle) and skin prick (blunt needle), using a single blind, randomised crossover design with 14 patients suffering from painful osteoarthritis, who were scanned with positron emission tomography (PET).⁷⁸ The results of the PET scans showed that insula ipsilateral to the site of stimulus was activated to a greater extent during real acupuncture than during the 'placebo control' intervention. Both 'real' and 'placebo control' acupuncture with the same expectation of effect caused greater activation than skin prick (which had no expectation of a therapeutic effect) in the right dorsolateral prefrontal cortex, anterior cingulate cortex, and midbrain. These results suggest that 'real' acupuncture and 'placebo control' acupuncture have specific physiological effects and that patients' expectation and belief regarding a potentially beneficial treatment modulate activity in cortical and subcortical areas. That 'placebo control' acupuncture may have a physiological effect is also supported by recent studies showing that stimulation of skin mechanoreceptors coupled to unmyelinated afferent C nerve fibres result in activity in the insular region.² Activity in these C tactile afferents has been implicated in a limbic reward response. It is likely that control procedures used in many acupuncture studies (superficial or minimal needling) aimed at being

inert are in fact activating these C tactile afferents and consequently are not inert. Furthermore, these 'control procedures' probably activate the reward system and as such induce feelings of wellbeing.²

Default mode

It has been suggested that PCC, rostral ACC, and MPFC compose a network characterised by a default mode of high level baseline activity, and further, that this baseline is similar to the brain's continuous resting oxygen extraction level. Possibly the default mode encompasses processes similar to introspection and the time relaxing between and after an acupuncture treatment.^{57,79-82} This is presently under investigation by Kathleen Hui and collaborators.

Acupuncture, reward and neuromodulation

Limbic structures play a key role in most diseases and illness responses including chronic pain and depression, regulating mood and neuromodulatory responses (eg sensory, autonomic, and endocrine). The pleasurable and neuromodulatory aspects of acupuncture as well as 'placebo needling' may partly be explained by the activation or deactivation of limbic structures including the hippocampus, amygdala, and their connections with the hypothalamus, septal area, and portions of the tegmentum.⁸³ As the limbic structures are activated or deactivated following stimulation of receptors in the skin, disregarding site, 'placebo-needling' does not exist and conclusions drawn on the basis that it is an inert control are false and truly misleading.

The hippocampus and frontal regions of the cerebral cortex have received much attention since reductions in hippocampal (and cortical) volumes were reported in patients with chronic pain, depression and post-traumatic stress disorder in whom memory and cognitive dysfunctions are commonly reported. Also, a decline in hippocampal function, which exerts inhibitory control over the hypothalamic-pituitary-adrenal (HPA) axis, could contribute to the hypercortisolemia found in a subset of depressed individuals. Bucinskaite and co-workers previously reported on the effects of repeated electroacupuncture treatments on open-field behaviour and on hippocampus concentrations of neuropeptide Y (NPY), neurokinin A (NKA), substance P (SP), galanin (GAL) and vasoactive intestinal peptide (VIP)-like immunoreactivities (-LI)

in the two rat strains Wistar-Kyoto (WKY) and spontaneously hypertensive rats (SHR).⁸⁴ Significantly higher concentrations of SP-LI, NKA-LI and NPY-LI were found in the hippocampus immediately after three weeks of electroacupuncture treatment compared to untreated animals with similar changes in neuropeptide concentrations in the two rat strains. Open-field behaviour was significantly reduced during the treatment period in both strains. Also, rats receiving treatment had higher thresholds to nociceptive stimuli and were calmer than control rats. There were significant negative correlations between behaviour and neuropeptide concentrations in spontaneously hypertensive rats, suggesting interdependency with sympathetic activity. It was proposed that some of the effects of electroacupuncture in rats are related to increases in NPY-LI, NKA-LI and SP-LI in the hippocampus. Interestingly, it has been reported that basal NPY-LI is lower in the hippocampus of 'depressed' rats (Flinders Sensitive Line) compared with controls, and that electroconvulsive stimuli (ECS) raise NPY-LI in the hippocampus.⁸⁵ These findings suggest that NPY is involved in depressive disorder and that antidepressant effects of ECS may in part be mediated through NPY. Furthermore, the hippocampus has been implicated in the regulation of anxiety and memory processes and low concentrations of SP-LI and NKA-LI has been seen in rats with chronic 'pain'. Possibly, this disturbance could be compensated for by repeated electroacupuncture treatments. Another, modality of stimulation that may prove to be effective in depression is laser-acupuncture (Quah-Smith, personal communication).

While the hippocampus and frontal cortex are involved in aspects of chronic pain and mood disorders, these regions do not account for all symptoms.⁸⁶⁻¹⁰¹ In recent years a role for other parts of the brain including the brain's reward regions have been demonstrated. Studies have identified the nucleus accumbens (NAcc) and its dopaminergic inputs from the ventral tegmental area (VTA) of the midbrain, as one of the most important anatomical substrates for rewards, such as food, sex, and social interactions. The amygdala, traditionally viewed as being critical for learned associations between negative emotional stimuli and environmental cues, serves a similar function for rewarding stimuli.

Interestingly, many patients with chronic pain and depression show reduced ability to experience pleasure (anhedonia) and loss of motivation, as well as abnormalities in several neurovegetative functions such as appetite, sleep, energy level, and circadian rhythms. All of these processes are part of highly overlapping and interacting circuits in which the dopaminergic VTA-NAcc pathway plays a critical role.¹⁰²⁻¹³⁵ A role for acupuncture and moxibustion in these conditions is supported by studies showing that they result in the activation of the reward system.⁷⁸

Probably, the transcription factor CREB (cAMP response element binding protein) is a key regulator of the reactivity of brain reward circuits and thereby regulates individual sensitivity to emotional stimuli.¹³⁵⁻¹⁴⁸ Short term increases in CREB activity in NAcc, induced by normal rewarding or aversive stimuli, could serve to dampen responses to subsequent stimuli and facilitate the ability to deal actively with the situation at hand (eg consumption of reward, escape from danger) ie increased resilience. Under more pathological conditions, however, larger and more sustained increases in CREB activity, induced by drug abuse or excessive stress, would lead to an excessive dampening of emotional reactivity. Conversely, sustained reductions in CREB activity, which are seen under conditions of social isolation, would heighten emotional reactivity and in the extreme be associated with a state of anxiety. The regulation of depression-like behaviour by changes in CREB activity within the NAcc is partly mediated by dynorphin.¹⁴⁹⁻¹⁵² Possibly, excessive activation of CREB by chronic stress increases dynorphin expression in the NAcc, which feed back to decrease VTA dopamine function and trigger certain features of chronic pain and depression. Recent findings indicate that acupuncture as well as moxibustion like stimulation modulates CREB (unpublished observations).

Brain derived nerve growth factor (BDNF) and other neurotrophins like nerve growth factor (NGF) play an important role in the reward system.¹⁵³⁻¹⁶² Under normal conditions, this BDNF signalling is critical for the appropriate memories of potentially rewarding or dangerous settings. Under pathological conditions, however, this signalling may establish abnormal associations, which would lead to certain symptoms of pain and depression even in the absence

of true external threats. Probably, BDNF signalling in the VTA-NAcc is required for the establishment of important associations with negative emotional stimuli. This could possibly explain why some patients who have tried acupuncture and experienced it as being unpleasant are less inclined to try it again as the mere thought could evoke unpleasant associations.

The hypothalamus plays an important role in reward mechanisms and in the neuromodulatory responses (sensory, autonomic and endocrine) to illness.¹⁶³⁻¹⁷² This has been explained in part by the fact that dopaminergic fibres from the VTA project to the NAcc through the lateral hypothalamus. Studies of feeding behaviour have especially provided new insights into the interaction between hypothalamus and VTA-NAcc. Recent work has begun to draw connections between hypothalamic feeding peptides and depression. Of particular interest in this respect is melanin-concentrating hormone (MCH), which is a major orexigenic (pro-appetite) peptide expressed in a subset of lateral hypothalamic neurons. Administration of MCH into the NAcc stimulates feeding behaviour, whereas blockade of its receptor (MCH1 receptor) decreases feeding. Moreover, several MCH1 receptor antagonists administered systemically or directly into the NAcc, exert antidepressant-like effects. Another transmitter in the lateral hypothalamus that also plays a role is orexin (hypocretin). Orexin increases feeding by promoting a state of wakefulness and alertness via orexin OX1 receptors. Further, orexin is also involved in regulating sleep-wake cycles. Abnormalities in orexin signalling have been implicated in disparate types of sleep abnormalities reported in chronic pain and depression. Taken together, two types of patients emerge: one whose chronic pain and depression is characterised by reduced activity and weight gain and one who exhibits increased activity, anxiety, and weight loss. An anti-depressant acting on one subset of receptors may therefore be effective or ineffective whereas acupuncture may work differently 're-establishing the balance' between the regulating centres disregarding the 'subtype'.

Many patients report their most serious symptoms in the morning with some improvement as the day progresses. This may represent an imbalance in their circadian rhythms causing fluctuations in mood, motivation, energy level, and responses to rewarding

stimuli.¹⁷³⁻¹⁷⁶ The suprachiasmatic nucleus (SCN) of the hypothalamus is considered the master circadian pacemaker of the brain. Here, circadian rhythms are generated at the molecular level by different transcription factors like Clock, Bmal, Per and Cry genes. Clock dimerises with Bmal and that dimer induces the expression of Per and Cry genes, which in turn feedback to repress Clock-Bmal activity. This Clock-Per cycle in the SCN is essential for matching circadian rhythms with the light-dark cycle. These circadian genes are also found in the VTA and NAcc indicating that they play a role in the regulation of mood. Furthermore, there has been interest in NPAS2 (neuronal Pas domain protein-2) in mood regulation.¹⁷⁷⁻¹⁷⁹ NPAS2 is homologous to Clock and, like Clock, dimerises with Bmal to regulate the expression of Per and Cry in a circadian fashion. NPAS2 is not expressed in SCN but in several limbic structures and especially within the NAcc.²⁰ Interestingly, NPAS2 knockout mice show increased anxiety-like behaviour and deficits in fear conditioning. It has been suggested that NPAS2 is a critical mediator of circadian rhythms in emotional responses via actions in limbic regions of the brain such as the NAcc. Preliminary findings suggest that acupuncture regulates NPAS2 activity.

Further studies are needed to explore the basis of acupuncture treatment in reward and reinforcement (delayed or uncertain), in implicit or procedural (stimulus-response) representational systems and in explicit or declarative representational systems. Individual differences in sensitivity to delays and uncertainty may contribute to the variable responses to acupuncture. Learning (attributing a meaning) and choice (preference) with delayed and uncertain reinforcement are related but in some cases dissociable processes. The contributions to 'delay discounting' and 'uncertainty discounting' of neuromodulators including serotonin and dopamine, and of specific neural structures have recently been reviewed.¹⁸⁰ Activation of the reward system by acupuncture stimulation is strongly supported by recent studies investigating the effects of electroacupuncture on serotonin and dopamine contents in the brain rewarding system in restrained conscious rats.¹⁸¹ The results showed that electroacupuncture increased the serotonin and dopamine contents of the nuclei accumbens, caudate, putamen and lateral hypothalamus, whereas

decreased content of these monoamines were seen in the dorsal raphe nucleus and amygdala. Differences in frequency effects (1Hz or 100Hz) on dopamine and serotonin were also reported. It was indicated that the effects of electroacupuncture compensated for the changes elicited by the restraint stress.

Acupuncture and long term effects

Napadow, Hui and collaborators recently used functional magnetic resonance imaging (fMRI) to evaluate the short and long term effects of acupuncture and 'sham' stimulation at acupoint LI4 in patients with carpal tunnel syndrome and in healthy controls.¹⁸⁶ Carpal tunnel syndrome patients responded to acupuncture with greater activation in the hypothalamus and deactivation in the amygdala as compared with healthy controls. A similar difference was found between carpal tunnel syndrome patients at baseline and after acupuncture treatments. For baseline carpal tunnel syndrome patients responding to acupuncture, functional connectivity was found between the hypothalamus and amygdala – the less the deactivation in the amygdala, the greater the activation in the hypothalamus, and vice versa. Furthermore, hypothalamic response correlated positively with the degree of maladaptive cortical plasticity in patients with carpal tunnel syndrome. This study provides evidence that chronic pain patients have a different response to acupuncture than healthy controls, through a coordinated network including the hypothalamus and amygdala. One reason that some previous studies on acupuncture imaging have not reported on the deactivation effects of acupuncture is possibly related to the fact that some authors, who use Statistical Parametric Mapping (SPM) for data analysis, have refrained from reporting deactivation, even if present, because of the dispute in the interpretation of blood oxygen level-dependant (BOLD) signal decreases.

In addition to the mentioned networks above, the cortical-cerebellar system needs to be taken into account. Ramnani and collaborators argue that a cortical-cerebellar system is central to the development of neural models that learn and eventually stimulate routinely-executed cognitive processes.^{184;185} Initially, higher order cognitive processes, mediated in prefrontal cortical loci, shift control iteratively to internal cerebellar representations of these processes, resulting in sparing

processing load on less efficient executive pre-frontal resources. Evidence exists for this cortical–cerebellar system in the acquisition of motor skills, which enables rapid 'effortless' execution and sequencing of bodily movements.^{43;186-192} Furthermore, a growing body of results indicate that internal cerebellar models may extend across a much broader range of contributory cognitive processing, inclusive of attention to anticipated informational cues, processing prediction errors for monetary reward, and autobiographical memory. As with highly efficient cerebellar models of motor actions, which simulate complex bodily actions from simple motor commands, so too these models may facilitate detection processes inherent to appraising self-relevant content from task features, or, attentional shifting between one's external environment and introspective mental content.^{43;186-192} The influences of task relevance to the self may determine the effects of acupuncture suggesting that not only the pain system should be taken into account when assessing the effects of acupuncture. Also, part of the long term healing effects of acupuncture may be ascribed to changes in the cerebellar system thereby sparing processing load in cortical and subcortical areas.

Conclusion

In conclusion, the results of current research suggest that acupuncture techniques as well as non-penetrating 'placebo controls' activate the patients' expectation and belief regarding a potentially beneficial treatment thereby modulating activity in the reward system.^{193;194} Acupuncture may be viewed as an intervention that triggers and enhances expectancy but not equivalent with expectancy itself. These mechanisms are activated in addition to the well known analgesic effects of acupuncture.^{195;196}

'Self' may be seen as a shifting illusion, ceaselessly constructed and deconstructed, and the effect of acupuncture may reflect its status (as well as that of the therapist).

Further studies on acupuncture and the default mode are essential when assessing the effects of acupuncture using the neuroimaging paradigm in general.

Reference list

1. Andersson S, Lundeberg T. Acupuncture – from empiricism to science: functional background to acupuncture effects in pain and disease. *Med Hypotheses* 1995;45(3):271-81.

2. Lund I, Lundeberg T. Are minimal, superficial, acupuncture or sham acupuncture procedures acceptable as inert placebo controls? *Acupunct Med* 2006;24(1):13-5.
3. Thomas M, Lundeberg T. Does acupuncture work? *Pain Clinical Updates* 1996;IV(3):1-8.
4. Naslund J, Naslund UB, Odenbring S, Lundeberg T. Comparison of symptoms and clinical findings in subgroups of individuals with patellofemoral pain. *Physiother Theory Pract* 2006;22(3):105-18.
5. Naslund J, Naslund UB, Odenbring S, Lundeberg T. Sensory stimulation (acupuncture) for the treatment of idiopathic anterior knee pain. *J Rehabil Med* 2002; 34(5):231-8.
6. Hui KKS, Liu J, Marina O, Napadow V, Haselgrove C, Kwong KK, Kennedy DN, Makris N. The integrated response of the human cerebro-cerebellar and limbic systems to acupuncture stimulation at ST36 as evidenced by fMRI. *NeuroImage* 2005;27(3):479-96.
7. Wu MT, Sheen JM, Chuang KH, Yang P, Chin SL, Tsai CY, Chen CJ, Liao JR, Lai PH, Chu KA, Pan HB, Yang CF. Neuronal specificity of acupuncture response: a fMRI study with electroacupuncture. *Neuroimage* 2002; 16(4):1028-37.
8. Kawakita K, Shinbara H, Imai K, Fukuda F, Yano T, Kuriyama K. How do acupuncture and moxibustion act? Focusing on the progress in Japanese acupuncture research. *J Pharmacol Sci* 2006;100(5):443-59.
9. Schmitz TW, Johnson SC. Relevance to self: A brief review and framework of neural systems underlying appraisal. *Neurosci Biobehav Rev* 2007;31(4):585-96.
10. Amodio DM, Frith CD. Meeting of minds: the medial frontal cortex and social cognition. *Nat Rev Neurosci* 2006;7(4):268-77.
11. Anderson AK, Phelps EA. Lesions of the human amygdala impair enhanced perception of emotionally salient events. *Nature* 2001;411(6835):305-9.
12. Bartels A, Zeki S. The neural correlates of maternal and romantic love. *Neuroimage* 2004;21(3):1155-66.
13. Bornhoved K, Quante M, Glauche V, Bromm B, Weiller C, Buchel C. Painful stimuli evoke different stimulus-response functions in the amygdala, prefrontal, insula and somatosensory cortex: a single-trial fMRI study. *Brain* 2002;125(Pt 6):1326-36.
14. Craig AD. Forebrain emotional asymmetry: a neuro-anatomical basis? *Trends Cogn Sci* 2005;9(12): 566-71.
15. Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting interoceptive awareness. *Nat Neurosci* 2004;7(2):189-95.
16. Antoniadis EA, Winslow JT, Davis M, Amaral DG. 2007 Role of the primate amygdala in fear-potentiated startle: effects of chronic lesions in the rhesus monkey. *J Neurosci*. 2007; 11(28):7386-96
17. Devinsky O, Morrell MJ, Vogt BA. Contributions of anterior cingulate cortex to behaviour. *Brain* 1995;118(Pt 1):279-306.
18. Elliott R, Newman JL, Longe OA, Deakin JF. Differential response patterns in the striatum and orbitofrontal cortex to financial reward in humans: a parametric functional magnetic resonance imaging study. *J Neurosci* 2003; 23(1):303-7.
19. Hamann S, Mao H. Positive and negative emotional verbal stimuli elicit activity in the left amygdala. *NeuroReport* 2002;13(1):15-9.
20. Kim H, Somerville LH, Johnstone T, Polis S, Alexander AL, Shin LM, Whalen PJ. Contextual modulation of amygdala responsivity to surprised faces. *J Cogn Neurosci* 2004;16(10):1730-45.
21. Knutson B, Fong GW, Bennett SM, Adams CM, Hommer D. A region of mesial prefrontal cortex tracks monetarily rewarding outcomes: characterization with rapid event-related fMRI. *Neuroimage* 2003;18(2):263-72.
22. Knutson B, Taylor J, Kaufman M, Peterson R, Glover G. Distributed neural representation of expected value. *J Neurosci* 2005;25(19):4806-12.
23. Kulkarni B, Bentley DE, Elliott R, Youell P, Watson A, Derbyshire SW, Frackowiak RS, Friston KJ, Jones AK. Attention to pain localization and unpleasantness discriminates the functions of the medial and lateral pain systems. *Eur J Neurosci* 2005;21(11):3133-42.
24. LeDoux JE. Emotion circuits in the brain. *Annu Rev Neurosci* 2000;23:155-84.
25. Liberzon I, Phan KL, Decker LR, Taylor SF. Extended amygdala and emotional salience: a PET activation study of positive and negative affect. *Neuropsychopharmacology* 2003;28(4):726-33.
26. Morgane PJ, Galler JR, Mokler DJ. A review of systems and networks of the limbic forebrain/limbic midbrain. *Prog Neurobiol* 2005;75(2):143-60.
27. Morris JS, Dolan RJ. Dissociable amygdala and orbitofrontal responses during reversal fear conditioning. *Neuroimage* 2004;22(1):372-80.
28. Nitschke JB, Sarinopoulos I, Mackiewicz KL, Schaefer HS, Davidson RJ. Functional neuroanatomy of aversion and its anticipation. *Neuroimage* 2006;29(1):106-16.
29. Northoff G, Bermpohl F. Cortical midline structures and the self. *Trends Cogn Sci* 2004;8(3):102-7.
30. Northoff G, Heinzel A, de Greck M, Bermpohl F, Dobrowolny H, Panksepp J. Self-referential processing in our brain—a meta-analysis of imaging studies on the self. *Neuroimage* 2006;31(1):440-57.
31. Ochsner KN, Beer JS, Robertson ER, Cooper JC, Gabrieli JD, Kihlstrom JF, D'Esposito M. The neural correlates of direct and reflected self-knowledge. *Neuroimage* 2005; 28(4):797-814.
32. Pessoa L, McKenna M, Gutierrez E, Ungerleider LG. Neural processing of emotional faces requires attention. *Proc Natl Acad Sci U S A* 2002;99(17):11458-63.
33. Pessoa L, Padmala S, Morland T. Fate of unattended fearful faces in the amygdala is determined by both attentional resources and cognitive modulation. *Neuroimage* 2005; 28(1):249-55.
34. Phan KL, Wager T, Taylor SF, Liberzon I. Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *Neuroimage* 2002; 16(2):331-48.
35. Phan KL, Taylor SF, Welsh RC, Decker LR, Noll DC, Nichols TE, Britton JC, Liberzon I. Activation of the medial prefrontal cortex and extended amygdala by individual ratings of emotional arousal: a fMRI study. *Biol Psychiatry* 2003;53(3):211-5.
36. Phan KL, Taylor SF, Welsh RC, Ho SH, Britton JC, Liberzon I. Neural correlates of individual ratings of emotional salience: a trial-related fMRI study. *Neuroimage* 2004;21(2):768-80.
37. Phelps EA, Delgado MR, Nearing KI, LeDoux JE.

- Extinction learning in humans: role of the amygdala and vmPFC. *Neuron* 2004;43(6):897-905.
38. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biol Psychiatry* 2003;54(5):504-14.
 39. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biol Psychiatry* 2003;54(5):515-28.
 40. Salomons TV, Johnstone T, Backonja MM, Davidson RJ. Perceived controllability modulates the neural response to pain. *J Neurosci* 2004;24(32):7199-203.
 41. Schaefer A, Collette F, Philippot P, van der Linden M, Laureys S, Delfiore G, Degueldre C, Maquet P, Luxen A, Salmon E. Neural correlates of "hot" and "cold" emotional processing: a multilevel approach to the functional anatomy of emotion. *Neuroimage* 2003;18(4):938-49.
 42. Singer T, Seymour B, O'Doherty J, Kaube H, Dolan RJ, Frith CD. Empathy for pain involves the affective but not sensory components of pain. *Science* 2004;303(5661):1157-62.
 43. Svoboda E, McKinnon MC, Levine B. The functional neuroanatomy of autobiographical memory: a meta-analysis. *Neuropsychologia* 2006;44(12):2189-208.
 44. Wager TD, Rilling JK, Smith EE, Sokolik A, Casey KL, Davidson RJ, Kosslyn SM, Rose RM, Cohen JD. Placebo-induced changes in fMRI in the anticipation and experience of pain. *Science* 2004;303(5661):1162-7.
 45. Wagner AD, Shannon BJ, Kahn I, Buckner RL. Parietal lobe contributions to episodic memory retrieval. *Trends Cogn Sci* 2005;9(9):445-53.
 46. Whalen PJ, Rauch SL, Etcoff NL, McInerney SC, Lee MB, Jenike MA. Masked presentations of emotional facial expressions modulate amygdala activity without explicit knowledge. *J Neurosci* 1998;18(1):411-8.
 47. Whalen PJ, Shin LM, Somerville LH, McLean AA, Kim H. Functional neuroimaging studies of the amygdala in depression. *Semin Clin Neuropsychiatry* 2002;7(4):234-42.
 48. Winston JS, Gottfried JA, Kilner JM, Dolan RJ. Integrated neural representations of odor intensity and affective valence in human amygdala. *Neurosci* 2005;25(39):8903-7.
 49. Zink CF, Pagnoni G, Martin ME, Dhamala M, Berns GS. Human striatal response to salient nonrewarding stimuli. *J Neurosci* 2003;23(22):8092-7.
 50. Zink CF, Pagnoni G, Martin-Skurski ME, Chappelow JC, Berns GS. Human striatal responses to monetary reward depend on saliency. *Neuron* 2004;42(3):509-17.
 51. Burgess N, Maguire EA, Spiers HJ, O'Keefe J. A temporoparietal and prefrontal network for retrieving the spatial context of lifelike events. *Neuroimage* 2001;14(2):439-53.
 52. Craik FIM, Moroz TM, Moscovitch M, Stuss DT, Winocur G, Tulving E, Kapur S. In search of the self: A positron emission tomography study. *Psychol Sci* 1999;10(1):26-34.
 53. Cunningham WA, Johnson MK, Gatenby JC, Gore JC, Banaji MR. Neural components of social evaluation. *J Pers Soc Psychol* 2003;85(4):639-49.
 54. Cunningham WA, Raye CL, Johnson MK. Implicit and explicit evaluation: fMRI correlates of valence, emotional intensity, and control in the processing of attitudes. *J Cogn Neurosci* 2004;16(10):1717-29.
 55. Greene JD, Sommerville RB, Nystrom LE, Darley JM, Cohen JD. An fMRI investigation of emotional engagement in moral judgment. *Science* 2001;293(5537):2105-8.
 56. Greene JD, Nystrom LE, Engell AD, Darley JM, Cohen JD. The neural bases of cognitive conflict and control in moral judgment. *Neuron* 2004;44(2):389-400.
 57. Gusnard DA, Akbudak E, Shulman GL, Raichle ME. Medial prefrontal cortex and self-referential mental activity: relation to a default mode of brain function. *Proc Natl Acad Sci U S A* 2001;98(7):4259-64.
 58. Henson RN, Rugg MD, Shallice T, Josephs O, Dolan RJ. Recollection and familiarity in recognition memory: an event-related functional magnetic resonance imaging study. *J Neurosci* 1999;19(10):3962-72.
 59. Jacobsen T, Schubotz RI, Hofel L, Cramon DY. Brain correlates of aesthetic judgment of beauty. *Neuroimage* 2006;29(1):276-85.
 60. Johnson SC, Baxter LC, Wilder LS, Pipe JG, Heiserman JE, Prigatano GP. Neural correlates of self-reflection. *Brain* 2002;125(Pt 8):1808-14.
 61. Johnson SC, Schmitz TW, Kawahara-Baccus TN, Rowley HA, Alexander AL, Lee J, Davidson RJ. The cerebral response during subjective choice with and without self-reference. *J Cogn Neurosci* 2005;17(12):1897-906.
 62. Kelley WM, Macrae CN, Wyland CL, Caglar S, Inati S, Heatherton TF. Finding the self? An event-related fMRI study. *J Cogn Neurosci* 2002;14(5):785-94.
 63. Maddock RJ. The retrosplenial cortex and emotion: new insights from functional neuroimaging of the human brain. *Trends Neurosci* 1999;22(7):310-6.
 64. Moll J, Eslinger PJ, Oliveira-Souza R. Frontopolar and anterior temporal cortex activation in a moral judgment task: preliminary functional MRI results in normal subjects. *Arq Neuropsiquiatr* 2001;59(3-B):657-64.
 65. Moll J, de Oliveira-Souza R, Eslinger PJ, Bramati IE, Mourão-Miranda J, Andreiuolo PA, Pessoa L. The neural correlates of moral sensitivity: a functional magnetic resonance imaging investigation of basic and moral emotions. *J Neurosci* 2002;22(7):2730-6.
 66. Nielsen FA, Balslev D, Hansen LK. Mining the posterior cingulate: segregation between memory and pain components. *Neuroimage* 2005;27(3):520-32.
 67. Piefke M, Weiss PH, Zilles K, Markowitsch HJ, Fink GR. Differential remoteness and emotional tone modulate the neural correlates of autobiographical memory. *Brain* 2003;126(Pt 3):650-68.
 68. Schmitz TW, Johnson SC. Self-appraisal decisions evoke dissociated dorsal-ventral aMPFC networks. *Neuroimage* 2006;30(3):1050-8.
 69. Schmitz TW, Kawahara-Baccus TN, Johnson SC. Metacognitive evaluation, self-relevance, and the right prefrontal cortex. *Neuroimage* 2004;22(2):941-7.
 70. Seger CA, Stone M, Keenan JP. Cortical Activations during judgments about the self and an other person. *Neuropsychologia* 2004;42(9):1168-77.
 71. Shah NJ, Marshall JC, Zafiris O, Schwab A, Zilles K, Markowitsch HJ, Fink GR. The neural correlates of person familiarity: A functional magnetic resonance imaging study with clinical implications. *Brain* 2001;124(Pt 4):804-15.
 72. Voegeley K, May M, Ritzl A, Falkai P, Zilles K, Fink GR. Neural correlates of first-person perspective as one constituent of human self-consciousness. *J Cogn Neurosci* 2004;16(5):817-27.

73. Vogt BA, Finch DM, Olson CR. Functional heterogeneity in cingulate cortex: the anterior executive and posterior evaluative regions. *Cereb Cortex* 1992;2(6):435-43.
74. Vogt BA, Berger GR, Derbyshire SW. Structural and functional dichotomy of human midcingulate cortex. *Eur J Neurosci* 2003;18(11):3134-44.
75. Vogt BA, Vogt L, Laureys S. Cytology and functionally correlated circuits of human posterior cingulate areas. *Neuroimage* 2006;29(2):452-66.
76. Zysset S, Huber O, Ferstl E, von Cramon DY. The Anterior Frontomedian Cortex and Evaluative Judgment: An fMRI Study. *Neuroimage* 2002;15(4):983-91.
77. Zysset S, Huber O, Samson A, Ferstl EC, von Cramon DY. Functional specialization within the anterior medial prefrontal cortex: a functional magnetic resonance imaging study with human subjects. *Neurosci Lett* 2003;335(3):183-6.
78. Pariente J, White P, Frackowiak RS, Lewith G. Expectancy and belief modulate the neuronal substrates of pain treated by acupuncture. *Neuroimage* 2005;25(4):1161-7.
79. Fransson P. Spontaneous low-frequency BOLD signal fluctuations: an fMRI investigation of the resting-state default mode of brain function hypothesis. *Hum Brain Mapp* 2005;26(1):15-29.
80. Greicius MD, Menon V. Default-mode activity during a passive sensory task: uncoupled from deactivation but impacting activation. *J Cogn Neurosci* 2004;16(9):1484-92.
81. Greicius MD, Srivastava G, Reiss AL, Menon V. Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc Natl Acad Sci U S A* 2004;101(13):4637-42.
82. Lustig C, Snyder AZ, Bhakta M, O'Brien KC, McAvoy M, Raichle ME, Morris JC, Buckner RL. Functional deactivations: change with age and dementia of the Alzheimer type. *Proc Natl Acad Sci U S A* 2003;100(24):14504-9.
83. Lundeberg T. Some of the effects of acupuncture in knee pain may be due to activation of the reward system. *Acupunct Med* 2006;24 Suppl:S67-70.
84. Bucinskaite V, Theodorsson E, Crumpton K, Stenfors C, Ekblom, A, Lundeberg T. Effects of repeated sensory stimulation (electroacupuncture) and physical exercise (running) on open- field behaviour and concentrations of neuropeptides in the hippocampus in WKY and SHR rats. *Eur J Neurosci* 1996;8(2):382-7.
85. Jimenez-Vasquez PA, Diaz-Cabiale Z, Caberlotto L, Bellido I, Overstreet D, Fuxe K, Mathe AA. Electroconvulsive stimuli selectively affect behavior and neuropeptide Y (NPY) and NPY Y(1) receptor gene expressions in hippocampus and hypothalamus of Flinders Sensitive Line rat model of depression. *Eur Neuropsychopharmacol* 2007;17(4):298-308.
86. Brody AL, Barsom MW, Bota RG, Saxena S. Prefrontal-subcortical and limbic circuit mediation of major depressive disorder. *Semin Clin Neuropsychiatry* 2001;6(2):102-12.
87. Drevets WC. Neuroimaging and neuropathological studies of depression: implications for the cognitive-emotional features of mood disorders. *Curr Opin Neurobiol* 2001;11(2):240-9.
88. Drevets WC, Gautier C, Price JC, Kupfer DJ, Kinahan PE, Grace AA, Price JL, Mathis CA. Amphetamine-induced dopamine release in human ventral striatum correlates with euphoria. *Biol Psychiatry* 2001;49(2):81-96.
89. Duman RS, Heninger GR, Nestler EJ. A molecular and cellular hypothesis of depression. *Arch Gen Psychiatry* 1997;54(7):597-606.
90. Karolewicz B, Szebeni K, Stockmeier CA, Konick L, Overholser JC, Jurjus G, Roth BL, Ordway GA. Low nNOS protein in the locus coeruleus in major depression. *J Neurochem* 2004;91(5):1057-66.
91. Klimek V, Schenck JE, Han H, Stockmeier CA, Ordway GA. Dopaminergic abnormalities in amygdaloid nuclei in major depression: a postmortem study. *Biol Psychiatry* 2002;52(7):740-8.
92. Manji HK, Drevets WC, Charney DS. The cellular neurobiology of depression. *Nat Med* 2001;7(5):541-7.
93. Mayberg HS. Positron emission tomography imaging in depression: a neural systems perspective. *Neuroimaging Clin N Am* 2003;13(4):805-15.
94. Mayberg HS, Lozano AM, Voon V, McNeely HE, Seminowicz D, Hamani C, Schwab JM, Kennedy SH. Deep brain stimulation for treatment-resistant depression. *Neuron* 2005;45(5):651-60.
95. Morilak DA, Frazer A. Antidepressants and brain monoaminergic systems: a dimensional approach to understanding their behavioural effects in depression and anxiety disorders. *Int J Neuropsychopharmacol* 2004;7(2):193-218.
96. Muller MB, Holsboer F. Mice with mutations in the HPA-system as models for symptoms of depression. *Biol Psychiatry* 2006;59(12):1104-15.
97. Nestler EJ. Molecular basis of neural plasticity underlying addiction. *Nat Rev Neurosci* 2001;2(2):119-28.
98. Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM. Neurobiology of depression. *Neuron* 2002;34(1):13-25.
99. Rajkowska G. Depression: what we can learn from postmortem studies. *Neuroscientist* 2003;9(4):273-84.
100. Ressler KJ, Nemeroff CB. Role of serotonergic and noradrenergic systems in the pathophysiology of depression and anxiety disorders. *Depress Anxiety* 2000;12 Suppl 1:2-19.
101. Tremblay LK, Naranjo CA, Graham SJ, Herrmann N, Mayberg HS, Hevenor S, Busto UE. Functional neuroanatomical substrates of altered reward processing in major depressive disorder revealed by a dopaminergic probe. *Arch Gen Psychiatry* 2005;62(11):1228-36.
102. Baldo BA, Gual-Bonilla L, Sijapati K, Daniel RA, Landry CF, Kelley AE. Activation of a subpopulation of orexin/hypocretin-containing hypothalamic neurons by GABAA receptor-mediated inhibition of the nucleus accumbens shell, but not by exposure to a novel environment. *Eur J Neurosci* 2004;19(2):376-86.
103. Cyr M, Morissette M, Barden N, Beaulieu S, Rochford J, Di Paolo T. Dopaminergic activity in transgenic mice underexpressing glucocorticoid receptors: effect of antidepressants. *Neuroscience* 2001;102(1):151-8.
104. Davis M, Whalen PJ. The amygdala: vigilance and emotion. *Mol Psychiatry* 2001;6(1):13-34.
105. Di Chiara G, Loddo P, Tanda G. Reciprocal changes in prefrontal and limbic dopamine responsiveness to adverse and rewarding stimuli after chronic mild stress: implications for the psychobiology of depression. *Biol Psychiatry* 1999;46(12):1624-33.
106. Espejo EF, Miñano FJ. Prefrontocortical dopamine depletion

- induces antidepressant-like effects in rats and alters the profile of desipramine during Porsolt's test. *Neuroscience* 1999;88(2):609-15.
107. Everitt BJ, Cardinal RN, Parkinson JA, Robbins TW. Appetitive behavior: impact of amygdala-dependent mechanisms of emotional learning. *Ann N Y Acad Sci* 2003;985:233-50.
 108. Everitt BJ, Wolf ME. Psychomotor stimulant addiction: a neural systems perspective. *J Neurosci* 2002;22(9):3312-20.
 109. Fulton S, Woodside B, Shizgal P. Modulation of brain reward circuitry by leptin. *Science* 2000;287(5450):125-8.
 110. Goldman-Rakic PS, Muly EC 3rd, Williams GV. D(1) receptors in prefrontal cells and circuits. *Brain Res Brain Res Rev* 2000;31(2-3):295-301.
 111. Heimer L, Zahm DS, Churchill L, Kalivas PW, Wohltmann C. Specificity in the projection patterns of accumbal core and shell in the rat. *Neuroscience* 1991;41(1):89-125.
 112. Horger BA, Iyasere CA, Berhow MT, Messer CJ, Nestler EJ, Taylor JR. Enhancement of locomotor activity and conditioned reward to cocaine by brain-derived neurotrophic factor. *J Neurosci* 1999;19(10):4110-22.
 113. Horger BA, Roth RH. The role of mesoprefrontal dopamine neurons in stress. *Crit Rev Neurobiol* 1996;10(3-4):395-418.
 114. Hyman SE, Malenka RC. Addiction and the brain: The neurobiology of compulsion and its persistence. *Nat Rev Neurosci* 2001;2(10):695-703.
 115. Jensen J, McIntosh AR, Crawley AP, Mikulis DJ, Remington G, Kapur S. Direct activation of the ventral striatum in anticipation of aversive stimuli. *Neuron* 2003;40(6):1251-7.
 116. Kalivas PW. Glutamate systems in cocaine addiction. *Curr Opin Pharmacol* 2004;4(1):23-9.
 117. Kelley AE, Berridge KC. The neuroscience of natural rewards: relevance to addictive drugs. *Neurosci* 2002;22(9):3306-11.
 118. Kiefer F, Wiedemann K. Neuroendocrine pathways of addictive behaviour. *Addict Biol* 2004;9(3-4):205-12.
 119. Koob GF, Le Moal M. Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology* 2001;24(2):97-129.
 120. Louilot A, Besson C. Specificity of amygdalo-striatal interactions in the involvement of mesencephalic dopaminergic neurons in affective perception. *Neuroscience* 2000;96(1):73-82.
 121. Nestler EJ. Molecular basis of neural plasticity underlying addiction. *Nature Rev Neurosci* 2001;2(2):119-28.
 122. Nicoullon A, Coquerel A. Dopamine: a key regulator to adapt action, emotion, motivation and cognition. *Curr Opin Neurol* 2003;16 Suppl 2:S3-9.
 123. Pallis E, Themos K, Spyraiki C. Chronic desipramine treatment selectively potentiates somatostatin-induced dopamine release in the nucleus accumbens. *Eur J Neurosci* 2001;14(4):763-7.
 124. Pezze MA, Feldon J. Mesolimbic dopaminergic pathways in fear conditioning. *Prog Neurobiol* 2004;74(5):301-20.
 125. Rada P, Moreno SA, Tucci S, Gonzalez LE, Harrison T, Chau DT, Hoebel BG, Hernandez L. Glutamate release in the nucleus accumbens is involved in behavioral depression during the Porsolt swim test. *Neuroscience* 2003;119(2):557-65.
 126. Renard CE, Fiocco AJ, Clenet F, Hascoet M, Bourin M. Is dopamine implicated in the antidepressant-like effects of selective serotonin re-uptake inhibitors in the mouse forced swimming test? *Psychopharmacology (Berl)* 2001;159(1):42-50.
 127. West AR, Floresco SB, Charara A, Rosenkranz JA, Grace AA. Electrophysiological interactions between striatal glutamatergic and dopaminergic systems. *Ann N Y Acad Sci* 2003;1003:53-74.
 128. West CH, Bonsall RW, Emery MS, Weiss JM. Rats selectively bred for high and low swim-test activity show differential responses to dopaminergic drugs. *Psychopharmacology (Berl)* 1999;146(3):241-51.
 129. Wise RA. Neuroleptics and operant behavior: The anhedonia hypothesis. *Behav Brain Sci* 1982;5:39-87.
 130. Wise RA. Addictive drugs and brain stimulation reward. *Annu Rev Neurosci* 1996;19:319-40.
 131. Wise RA. Drug-activation of brain reward pathways. *Drug Alcohol Depend* 1998;51(1-2):13-22.
 132. Wittmann BC, Schott BH, Guderian S, Frey JU, Heinze HJ, Duzel E. Reward-related fMRI activation of dopaminergic midbrain is associated with enhanced hippocampus-dependent long term memory formation. *Neuron* 2005;45(3):459-67.
 133. Yadid G, Overstreet DH, Zangen A. Limbic dopaminergic adaptation to a stressful stimulus in a rat model of depression. *Brain Res* 2001;896(1-2):43-7.
 134. Nestler EJ, Carlezon WA Jr. The mesolimbic dopamine reward circuit in depression. *Biol Psychiatry* 2006;59(12):1151-9.
 135. Barrot M, Olivier JD, Perrotti LI, DiLeone RJ, Berton O, Eisch AJ, Impey S, Storm DR, Neve RL, Yin JC, Zachariou V, Nestler EJ. CREB activity in the nucleus accumbens shell controls gating of behavioral responses to emotional stimuli. *Proc Natl Acad Sci U S A* 2002;99(17):11435-40.
 136. Barrot M, Wallace DL, Bolaños CA, Graham DL, Perrotti LI, Neve RL, Chambliss H, Yin JC, Nestler EJ. Regulation of anxiety and initiation of sexual anxiety by CREB in the nucleus accumbens. *Proc Natl Acad Sci U S A* 2005;102(23):8357-62.
 137. Carlezon WA Jr, Duman RS, Nestler EJ. The many faces of CREB. *Trends Neurosci* 2005;28(8):436-45.
 138. Carlezon WA Jr, Thome J, Olson VG, Lane-Ladd SB, Brodtkin ES, Hiroi N, Duman RS, Neve RL, Nestler EJ. Regulation of cocaine reward by CREB. *Science* 1998;282(5397):2272-5.
 139. Chen AC, Shirayama Y, Shin KH, Neve RL, Duman RS. Expression of the cAMP response element binding protein (CREB) in hippocampus produces an antidepressant effect. *Biol Psychiatry* 2001;49(9):753-62.
 140. Conti AC, Maas JW Jr, Muglia LM, Dave BA, Vogt SK, Tran TT, Rayhel EJ, Muglia LJ. Distinct regional and subcellular localization of adenylyl cyclases type 1 and 8 in mouse brain. *Neuroscience* 2007;146(2):713-29.
 141. Conti AC, Blendy JA. Regulation of antidepressant activity by cAMP response element binding proteins. *Mol Neurobiol* 2004;30(2):143-55.
 142. Conti AC, Cryan JF, Dalvi A, Lucki I, Blendy JA. cAMP response element-binding protein is essential for the upregulation of brain-derived neurotrophic factor transcription, but not the behavioral or endocrine responses to antidepressant drugs. *J Neurosci* 2002;22(8):3262-8.

143. McClung CA, Nestler EJ. Regulation of gene expression and cocaine reward by CREB and DeltaFosB. *Nat Neurosci* 2003;6(11):1208-15.
144. Olson VG, Zabetian CP, Bolanos CA, Edwards S, Barrot M, Eisch AJ, Hughes T, Self DW, Neve RL, Nestler EJ. Regulation of drug reward by CREB: Evidence for two functionally distinct subregions of the ventral tegmental area. *J Neurosci* 2005;25(23):5553-62.
145. Pandey SC, Roy A, Zhang H, Xu T. Partial deletion of the cAMP response element-binding protein gene promotes alcohol-drinking behaviors. *J Neurosci* 2004;24(21):5022-30.
146. Pliakas AM, Carlson RR, Neve RL, Konradi C, Nestler EJ, Carlezon WA. Altered responsiveness to cocaine and increased immobility in the forced swim test associated with elevated CREB expression in the nucleus accumbens. *J Neurosci* 2001;21(18):7397-403.
147. Ramos BP, Birnbaum SG, Lindenmayer I, Newton SS, Duman RS, Arnsten AF. Dysregulation of protein kinase a signaling in the aged prefrontal cortex: new strategy for treating age-related cognitive decline. *Neuron* 2003;40(4):835-45.
148. Valverde O, Mantamadiotis T, Torrecilla M, Ugedo L, Pineda J, Bleckmann S, Gass P, Kretz O, Mitchell JM, Schütz G, Maldonado R. Modulation of anxiety-like behavior and morphine dependence in CREB-deficient mice. *Neuropsychopharmacology* 2004;29(6):1122-33.
149. Walters CL, Kuo YC, Blendy JA. Differential distribution of CREB in the mesolimbic dopamine reward pathway. *J Neurochem* 2003;87(5):1237-44.
150. Li Y, van den Pol AN. Differential target-dependent actions of coexpressed inhibitory dynorphin and excitatory hypocretin/orexin neuropeptides. *J Neurosci*. 2006;26(50):13037-47.
151. Mague SD, Pliakas AM, Todtenkopf MS, Tomasiewicz HC, Zhang Y, Stevens WC Jr, Jones RM, Portoghese PS, Carlezon WA Jr. Antidepressant-like effects of kappa-opioid receptor antagonists in the forced swim test in rats. *J Pharmacol Exp Ther* 2003;305(1):323-30.
152. McLaughlin JP, Marton-Popovici M, Chavkin C. Kappa opioid receptor antagonism and prodynorphin gene disruption block stress-induced behavioral responses. *J Neurosci* 2003;23(13):5674-83.
153. Newton SS, Thome J, Wallace TL, Shirayama Y, Schlesinger L, Sakai N, Chen J, Neve R, Nestler EJ, Duman RS. Inhibition of CREB or dynorphin in the nucleus accumbens produces an antidepressant-like effect. *J Neurosci* 2002;22(24):10883-90.
154. Shippenberg TS, Rea W. Sensitization to the behavioral effects of cocaine: modulation by dynorphin and kappa-opioid receptor agonists. *Pharmacol Biochem Behav* 1997;57(3):449-55.
155. Todtenkopf MS, Marcus JF, Portoghese PS, Carlezon WA. Effects of kappa-opioid receptor ligands on intracranial self-stimulation in rats. *Psychopharmacology (Berl)* 2004;172(4):463-70.
156. Berhow MT, Russell DS, Terwilliger RZ, Beitner-Johnson D, Self DW, Lindsay RM, Nestler EJ. Influence of neurotrophic factors on morphine- and cocaine-induced biochemical changes in the mesolimbic dopamine system. *Neuroscience* 1995;68(4):969-79.
157. Berton O, McClung CA, Dileone RJ, Krishnan V, Renthal W, Russo SJ, Graham D, Tsankova NM, Bolanos CA, Rios M, Monteggia LM, Self DW, Nestler EJ. Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* 2006;311(5762):864-8.
158. Buwalda B, Kole MH, Veenema AH, Huininga M, de Boer SF, Korte SM, Koolhaas JM. Long term effects of social stress on brain and behavior: a focus on hippocampal functioning. *Neurosci Biobehav Rev* 2005;29(1):83-97.
159. Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 2006;59(12):1116-27.
160. Eisch AJ, Bolaños CA, de Wit J, Simonak RD, Pudiak CM, Barrot M, Verhaagen J, Nestler EJ. BDNF in the ventral midbrain-nucleus accumbens pathway: a role in depression. *Biol Psychiatry* 2003;54(10):994-1005.
161. Hall FS, Drgonova J, Goeb M, Uhl GR. Reduced behavioral effects of cocaine in heterozygous brain-derived neurotrophic factor (BDNF) knockout mice. *Neuropsychopharmacology* 2003;28(8):1485-90.
162. Horger BA, Iyasere CA, Berhow MT, Messer CJ, Nestler EJ, Taylor JR. Enhancement of locomotor activity and conditioned reward to cocaine by brain-derived neurotrophic factor. *J Neurosci* 1999;19(10):4110-22.
163. Lu L, Dempsey J, Liu SY, Bossert JM, Shaham Y. A single infusion of brain-derived neurotrophic factor into the ventral tegmental area induces long-lasting potentiation of cocaine seeking after withdrawal. *J Neurosci* 2004;24(7):1604-11.
164. Monteggia LM, Barrot M, Powell CM, Berton O, Galanis V, Gemelli T, Meuth S, Nagy A, Greene RW, Nestler EJ. Essential role of BDNF in adult hippocampal function. *Proc Natl Acad Sci U S A* 2004;101(29):10827-32.
165. Pierce RC, Bari AA. The role of neurotrophic factors in psychostimulant-induced behavioral and neuronal plasticity. *Rev Neurosci* 2001;12(2):95-110.
166. Turner CA, Akil H, Watson SJ, Evans SJ. The fibroblast growth factor system and mood disorders. *Biol Psychiatry* 2006;59(12):1128-35.
167. Borowsky B, Durkin MM, Ogozalek K, Marzabadi MR, DeLeon J, Lagu B, Heurich R, Lichtblau H, Shaposhnik Z, Daniewska I, Blackburn TP, Branchek TA, Gerald C, Vaysse PJ, Forray C. Antidepressant, anxiolytic and anorectic effects of a melanin-concentrating hormone-1 receptor antagonist. *Nat Med* 2002;8(8):825-30.
168. Chaki S, Hirota S, Funakoshi T, Suzuki Y, Suetake S, Okubo T, Ishii T, Nakazato A, Okuyama S. Anxiolytic-like and antidepressant-like activities of MCL0129 (1-[(S)-2-(4-fluorophenyl)-2-(4-isopropylpiperidin-1-yl)ethyl]-4-[4-(2-methoxy-naphthalen-1-yl)butyl]piperazine), a novel and potent nonpeptide antagonist of the melanocortin-4 receptor. *J Pharmacol Exp Ther* 2003;304(2):818-26.
169. Daniels E, King MA, Smith IE, Shneerson JM. Health-related quality of life in narcolepsy. *J Sleep Res* 2001;10(1):75-81.
170. Georgescu D, Sears RM, Hommel JD, Barrot M, Bolanos CA, Marsh DJ, Bednarek MA, Bibb JA, Maratos-Flier E, Nestler EJ, DiLeone RJ. The hypothalamic neuropeptide melanin-concentrating hormone acts in the nucleus accumbens to modulate feeding behavior and forced-swim performance. *J Neurosci* 2005;25(11):2933-40.
171. Heilig M. The NPY system in stress, anxiety and depression. *Neuropeptides* 2004;38(4):213-24.

172. Hsu R, Taylor JR, Newton SS, Alvaro JD, Haile C, Han G, Hruby VJ, Nestler EJ, Duman RS. Blockade of melanocortin transmission inhibits cocaine reward. *Eur J Neurosci* 2005;21(8):2233-42.
173. Korotkova TM, Sergeeva OA, Eriksson KS, Haas HL, Brown RE. Excitation of ventral tegmental area dopaminergic and nondopaminergic neurons by orexins/hypocretins. *J Neurosci*. 2003;23(1):7-11.
174. Mignot E. Sleep, sleep disorders and hypocretin (orexin). *Sleep Med* 2004;5 Suppl 1:S2-8.
175. Nakamura T, Uramura K, Nambu T, Yada T, Goto K, Yanagisawa M, Sakurai T. Orexin-induced hyperlocomotion and stereotypy are mediated by the dopaminergic system. *Brain Res* 2000;873(1):181-7.
176. Saito Y, Nothacker HP, Wang Z, Lin SH, Leslie F, Civelli O. Molecular characterization of the melanin-concentrating-hormone receptor. *Nature* 1999;400(6741):265-9.
177. Abarca C, Albrecht U, Spanagel R. Cocaine sensitization and reward are under the influence of circadian genes and rhythm. *Proc Natl Acad Sci U S A* 2002;99(13):9026-30.
178. Andretic R, Chaney S, Hirsh J. Requirement of circadian genes for Cocaine sensitization in *Drosophila*. *Science* 1999;285(5430):1066-8.
179. McClung CA, Sidiropoulou K, Vitaterna M, Takahashi JS, White FJ, Cooper DC, Nestler EJ. Regulation of dopaminergic transmission and cocaine reward by the Clock gene. *Proc Natl Acad Sci U S A* 2005;102(26):9377-81.
180. Okamura H, Yamaguchi S, Yagita K. Molecular machinery of the circadian clock in mammals. *Cell Tissue Res* 2002;309(1):47-56.
181. Dudley CA, Erbel-Sieler C, Estill SJ, Reick M, Franken P, Pitts S, McKnight SL. Altered patterns of sleep and behavioral adaptability in NPAS2-deficient mice. *Science* 2003;301(5631):379-83.
182. Garcia JA, Zhang D, Estill SJ, Michnoff C, Rutter J, Reick M, Scott K, Diaz-Arrastia R, McKnight SL. Impaired cued and contextual memory in NPAS2-deficient mice. *Science* 2000;288(5474):2226-30.
183. Reick M, Garcia JA, Dudley C, McKnight SL. NPAS2: an analog of clock operative in the mammalian forebrain. *Science* 2001;293(5529):506-9.
184. Cardinal RN. Neural systems implicated in delayed and probabilistic reinforcement. *Neural Netw* 2006;19(8):1277-301.
185. Yoshimoto K, Fukuda F, Hori M, Kato B, Kato H, Hattori H, Tokuda N, Kuriyama K, Yano T, Yasuhara M. Acupuncture stimulates the release of serotonin, but not dopamine, in the rat nucleus accumbens. *Tohoku J Exp Med* 2006;208(4):321-6.
186. Napadow V, Kettner N, Liu J, Li M, Kwong KK, Vangel M, Makris N, Audette J, Hui KK. Hypothalamus and amygdala response to acupuncture stimuli in carpal tunnel syndrome. *Pain* 2007;130(3):254-66.
187. Ramnani N, Elliott R, Athwal BS, Passingham RE. Prediction error for free monetary reward in the human prefrontal cortex. *Neuroimage* 2004;23(3):777-86.
188. Ramnani N, Behrens TE, Johansen-Berg H, Richter MC, Pinsk MA, Andersson JL, Rudebeck P, Ciccirelli O, Richter W, Thompson AJ, Gross CG, Robson MD, Kastner S, Matthews PM. The evolution of prefrontal inputs to the cortico-pontine system: diffusion imaging evidence from Macaque monkeys and humans. *Cereb Cortex* 2006;16(6):811-8.
189. Allen G, Buxton RB, Wong EC, Courchesne E. Attentional activation of the cerebellum independent of motor involvement. *Science* 1997;275(5308):1940-3.
190. Bastian AJ, Thach WT. Cerebellar outflow lesions: a comparison of movement deficits resulting from lesions at the levels of the cerebellum and thalamus. *Ann Neurol* 1995;38(6):881-92.
191. Fiez JA. Cerebellar contributions to cognition. *Neuron* 1996;16(1):13-5.
192. Goodkin HP, Keating JG, Martin TA, Thach WT. Preserved simple and impaired compound movement after infarction in the territory of the superior cerebellar artery. *Can J Neurol Sci* 1993;20 Suppl 3:S93-104.
193. Imamizu H, Miyauchi S, Tamada T, Sasaki Y, Takino R, Pütz B, Yoshioka T, Kawato M. Human cerebellar activity reflecting an acquired internal model of a new tool. *Nature* 2000;403(6766):192-5.
194. Krupa DJ, Thompson JK, Thompson RF. Localization of a memory trace in the mammalian brain. *Science* 1993;260(5110):989-91.
195. Lu X, Hikosaka O, Miyachi S. Role of monkey cerebellar nuclei in skill for sequential movement. *J Neurophysiol* 1998;79(5):2245-54.
196. Dhond RP, Kettner N, Napadow V. Do the neural correlates of acupuncture and placebo effects differ? *Pain* 2007;128(1-2):8-12.
197. Linde K, Witt CM, Streng A, Weidenhammer W, Wagenpfeil S, Brinkhaus B, Willich SN, Melchart D. The impact of patient expectations on outcomes in four randomized controlled trials of acupuncture in patients with chronic pain. *Pain* 2007;128(3):264-71.
198. Lundeberg T, Hurtig T, Lundeberg S, Thomas M. Long term results of acupuncture in chronic head and neck pain. *Pain Clinic* 1988;2(1):15-31.
199. Han JS. Acupuncture: neuropeptide release produced by electrical stimulation of different frequencies. *Trends Neurosci* 2003;26(1):17-22.