

Are reviews based on sham acupuncture procedures in fibromyalgia syndrome (FMS) valid?

Thomas Lundeberg & Iréne Lund

Thomas Lundeberg,
senior consultant
Rehabilitation Medicine
University Clinic
Danderyds Hospital
Stockholm, Sweden

Iréne Lund
lecturer
registered
physiotherapist
Department of
Physiology and
Pharmacology
Karolinska Institutet
Stockholm, Sweden

Correspondence:
Thomas Lundeberg

thomas.lundeberg@ds.se

Abstract

In recent reviews regarding the efficacy of acupuncture in fibromyalgia syndrome (FMS) it has been concluded that acupuncture has no specific effect since the control procedure (superficial needling and/or needling away from 'specific' points) had similar effects. These conclusions may be questioned since superficial needling and/or needling away from specific trigger points is not inert. Also, manual acupuncture or mild electroacupuncture (EA) may not be sufficient to activate the endogenous pain inhibiting system.

Patients with FMS suffer from allodynia, fatigue and muscle ache, which is partly explained by peripheral and central sensitisation. Sensitisation results in augmented and altered stimulus responses whereby light stimulation of the skin has as strong an effect as regular needling on the pain inhibitory system in FMS.

Central sensitisation in FMS is also associated with expanded receptive fields of central neurons resulting in a larger topographic distribution of the pain. This would suggest that control procedures using needling away from the 'specific site' might have as strong an effect as needling within the most painful area. Also, repeated nociceptive input from muscles (as obtained by *de qi*) results in expansion of receptive fields which in turn may result in activation of descending pain inhibition outside the stimulated myotome.

Sensitisation to pain, such as in FMS, may also be related to abnormalities in descending efferent pathways. As there is likely to be an imbalance between excitatory and inhibitory systems in FMS, stronger stimulation may therefore be needed to activate the descending pain inhibitory system. In studies using mild manual acupuncture or weak EA stimulation optimal pain inhibition may therefore not have been obtained.

When conducting studies on acupuncture, the clinical condition or syndrome needs to be taken into account and the control procedure designed accordingly.

Keywords

Acupuncture, fibromyalgia syndrome, pain, placebo, sensitisation, sham control.

Introduction

In recent reviews evaluating the efficacy of acupuncture in fibromyalgia syndrome (FMS) it has been concluded that acupuncture has no specific effects since the control procedure (superficial needling and/or needling away from the 'specific' points) has similar effects.¹⁻¹¹ A prerequisite for these conclusions is that the control procedure used is inert and without specific effects. FMS is a chronic syndrome characterised by diffuse or specific muscle, joint, or bone pain, fatigue, and a wide range of other symptoms. The term fibromyalgia, coined in 1976 to more accurately describe the

symptoms, originates from the Latin word *fibra*, meaning fibre, and the Greek words *myo*, meaning muscle, and *algos*, meaning pain. The defining symptoms of FMS are chronic, widespread pain and tenderness to light muscle touch and moderate to severe fatigue. Those affected may also experience heightened sensitivity of the skin. In patients with FMS non-noxious stimuli give rise to pain, allodynia, and noxious stimuli cause pain that lasts longer and has higher intensity than normally, hyperalgesia. FMS is also characterised by muscle ache, muscle spasms, weakness in the limbs, and 'nerve' pain.¹²⁻²²

Pain in FMS

Central sensitisation of sensory neurons and pain disinhibition may explicate parts of the phenomenon seen in FMS.^{13-19;23} Central sensitisation implies spontaneous nerve activity, expanded receptive fields of central neurons, resulting in a larger topographic distribution of the pain, and augmented stimulus responses such as abnormal temporal summation.^{19;24-27} Peripheral sensitisation of sensory neurons innervating deeper tissues may also play a role.^{25;28}

In FMS, sensitisation of sensory transmission neurons in the central nervous system (CNS), especially the wide-dynamic-range (WDR) neurons, becomes permanent. The WDR neurons change in structure, phenotype, function and biochemistry. Afferent A δ nerve fibres that normally transmit sensations like touch, pressure and vibration get qualities that are similar to those of the thin afferent C-fibres (pain afferents). This implies that touch or slight pressure becomes painful.^{24;25;27}

Abnormal temporal summation, or 'wind-up', is the phenomenon whereby after an initial painful stimulus, subsequent equal stimuli are perceived to be more intensely painful.²⁹ This magnified 'second pain', which occurs in everyone, is exaggerated in patients with FMS.²⁶ The receptor thought to be responsible for these phenomena is the N-methyl-D-aspartate (NMDA) receptor,^{30;31} which is found at the postsynaptic membrane in the dorsal horn of the spinal cord. These receptors are normally inactive and do not respond to initial acute stimuli. However, after repeated neuronal depolarisation, the receptors undergo activation. Experimental evidences from animal and human studies have shown that NMDA receptors are responsible for wind-up and central sensitisation. NMDA receptors also play a critical role in neuronal plasticity and long term potentiation.^{32;33} It is likely that the NMDA receptors contribute to the pain in FMS^{24;25} as the temporal summation is attenuated by antagonists to NMDA receptors.^{19;27;34-45} Interestingly, it has been reported that electroacupuncture modulates the expression of NMDA receptors in primary sensory neurons,⁴⁶ and regulates NMDA receptor NR1 subunit expression,⁴⁷ suggesting that part of the effects of acupuncture may be attributed to modulation of NMDA receptor activity.

Whether activation of glial cells is important for the pathogenesis of pain and allodynia in FMS is currently unknown.⁴⁸⁻⁵⁰ Glial cells, long thought to be metabolically inactive support cells in the nervous system, are now recognised as playing a substantial role in modulating pain signalling.⁴⁸ Glial cells and astrocytes are activated by stimuli that induce pain, such as nerve trauma, subcutaneous irritation, and intraperitoneal inflammation,⁴⁹ and by neurotransmitters involved in pain signalling.⁴⁷ In addition to having receptors for neurotransmitters, glial cells express receptors for bacteria and viruses,⁵¹⁻⁵⁵ which may explain why infection with neurotropic organisms, such as HIV, is frequently associated with FMS or chronic pain. Glial cells release many neuroactive substances on activation by painful stimuli, including nitric oxide, prostaglandins, leukotrienes, nerve growth factors, excitatory amino acids, and reactive oxygens.⁵⁵ Activated glial cells induce the release of substance P and other excitatory amino acids from primary afferent neurons in the spinal cord and enhance the excitability of pain transmission neurons. In addition, microglia and astrocytes release proinflammatory cytokines, such as interleukin-1, interleukin-6, and tumor necrosis factor.⁵⁴ Increased levels of cytokines have been found in patients with FMS implicating a role in the aetiology of pain.⁵⁶⁻⁵⁸ Blocking the actions of these cytokines prevents or reverses exaggerated pain states.⁵⁹ Also, acupuncture has been shown to inhibit glial activation and inflammatory events including hyperalgesia.^{60;61}

Pain assessment and gender differences

To allow for assessment of the central sensitisation and the effects of acupuncture a reliable clinically applicable procedure is needed. We suggest that one such procedure is the assessment of matched pain and pain threshold using electrocutaneous stimulation,⁶²⁻⁶⁵ though women's and men's assessment should preferably be evaluated separately.⁶⁶ This recommendation is based on findings from a recent study where gender differences in pain perception were assessed following acupuncture. This study aimed to investigate changes in electrical sensory thresholds and electrical pain thresholds, in response to electroacupuncture (EA) in healthy women and men.⁶⁷ The thresholds were assessed pre-, during-, and post-EA. Equal levels of

systematic changes (changes in common for the group) towards increased electrical sensory thresholds were seen in women and men post-EA. At the same point of time, systematic changes towards increased electrical pain thresholds were only seen in women, while they were unchanged in men. Significant individual variations were found in the women's responses of assessed electrical sensory and pain thresholds but not in the men's. It is concluded that both women and men responded with a significant increase of the electrical sensory threshold to EA, but only women responded with increase of the electrical pain thresholds though with individual variation.

A great individual variation in response to acupuncture suggests that it is not possible to make a prediction about expected effects for the group and that the treatment should be tailored individually. From an individual patient perspective the only true evaluation of a treatment is to try it out (presupposed a safe treatment with limited side-effects). This suggestion is valid even if randomised controlled trials (RCTs) have concluded that there are limited effects of a treatment, since this conclusion often is based on the results of the group, ignoring the individual effects. These and other findings indicate that in evaluation of acupuncture treatment gender differences should be taken into account as well as the individual variation.

Control and treatment parameters

Superficial needling versus deep needling

Patients with FMS suffer from allodynia, fatigue and muscle ache partly explained by peripheral and central sensitisation. Sensitisation results in augmented and altered stimulus responses whereby light stimulation of the skin might have as strong an effect as regular needling on the pain inhibitory system in FMS, despite being used as an 'inert control'. This suggestion is supported by studies measuring blood flow in patients with FMS where superficial needling of the skin resulted in as strong an increase in blood flow as stimulation of muscle tissue. In healthy subjects superficial needling had no effect on blood flow.^{68:69}

Needling at specific points versus 'placebo' points

Recordings from neurons in the spinal dorsal horn have revealed that within minutes after application

of a noxious stimulus to a receptive field in a muscle, new receptive fields at a distance from the initial receptive field emerge.⁷⁰⁻⁷⁹ That is, following nociceptive input, dorsal horn neurons that were previously responsive to only one area in a muscle begin to respond to nociception from areas that previously had not provoked a response.^{28:71} This may be attributed to activation of NMDA receptors on WDR neurons causing additional release of neuropeptides such as substance P, and these substances can diffuse in the spinal cord and result in the spread of pain.^{20:37:42:57} Also, because of the several connections that exist between groups of glia and the types of transmitters they release, activation of glial cells may cause expansion of the receptive field or extraterritorial pain.^{48:49} This central sensitisation could explain the large topographic distribution and the referred pain seen in FMS. This would suggest that control procedures using needling away from the 'specific site' might have as strong an effect as needling within the most painful area.

Also, repeated nociceptive input from muscles (as obtained by *de qi*) results in expansion of receptive fields, which in turn may result in activation of descending pain inhibition outside the stimulated myotome, ie if one uses, 'non-specific' ('placebo') points outside the affected myotome (most painful area/s) there may still be an effect.^{80:81}

Altogether this would suggest that 'non-specific' points or sites do not exist in FMS. This may be true in other conditions as well.

Modality of stimulation

The endogenous pain modulation in FMS is impaired as compared to healthy controls.⁸²⁻⁸⁵ This disinhibition of pain is a result of a change in the endogenous descending pain-modulating system. The descending system includes a nervous network that links the periaqueductal grey and the rostral ventromedial medulla with the spinal cord.⁸⁶ In persons with permanent pain hypersensitivity there could be a continuous stimulation of both facilitatory and inhibitory tracts that would result in increased pain if activity in inhibitory tracts decreases or if activity in facilitatory tracts increases.⁸⁷ The descending facilitatory pathways originate in the frontal cortical areas,⁸⁸ and have been shown to cause generalised increased neuronal responses

along the neuroaxis, indicating that emotions such as fear may drive the development of widespread pain and sensitisation.²⁴⁻²⁵

As there probably is an imbalance between excitatory and inhibitory systems in FMS, stronger stimulation may therefore be needed to activate the descending pain inhibitory system. In studies using mild manual acupuncture or weak EA stimulation, optimal pain inhibition may therefore not have been obtained. This suggestion is supported by a study showing that strong electroacupuncture stimulation was needed to obtain pain alleviation in FMS.⁸⁹

However, clinical experiences indicate that some FMS patients experience more pain when treated with acupuncture. This would indicate that depending on the status of the pain inhibitory system two responses may be obtained. When there is a light to moderate dysfunction of endogenous pain inhibition, strong stimulation may prove to be effective and when there is a severe dysfunction no pain inhibition is obtained and instead the treatment results in more pain. This would suggest that both 'light' and 'strong' stimulation should be tried out. The patient should then be allowed to continue with the stimulation modality preferred, thereby also optimising expectancy.⁹⁰⁻⁹¹

Future perspective

In many randomised control trials a treatment is compared with its inactive control. However, a prerequisite is that one can design an appropriate control procedure. This may not be possible when assessing the effects of procedures like acupuncture. We therefore suggest that treatment procedures (for example 10 treatments by an acupuncturist versus a visit to a physician prescribing a drug which is then used) should be compared instead of separate treatments. Also, the efficacy of acupuncture should be compared with the recommended drug and its placebo taking side-effects, compliance and costs into account. In the future, environmental aspects of drugs (production, degradation etc) should be taken into account for the evaluation of optimal treatment strategies.⁹²⁻¹⁰³

Concluding remark

Taken together this would suggest that when conducting studies on acupuncture one needs to take the clinical condition or syndrome into account and design the control procedure appropriately.

Reference list

1. Mannerkorpi K, Henriksson C. Non-pharmacological treatment of chronic widespread musculoskeletal pain. *Best Pract Res Clin Rheumatol* 2007;21(3):513-34.
2. Rooks DS. Fibromyalgia treatment update. *Curr Opin Rheumatol* 2007;19(2):111-7.
3. Mayhew E, Ernst E. Acupuncture for fibromyalgia—a systematic review of randomized clinical trials. *Rheumatology (Oxford)* 2007;46(5):801-4.
4. Gur A. Physical therapy modalities in management of fibromyalgia. *Curr Pharm Des* 2006;12(1):29-35.
5. Zheng L, Faber K. Review of the Chinese medical approach to the management of fibromyalgia. *Curr Pain Headache Rep* 2005;9(5):307-12.
6. Ernst E. Musculoskeletal conditions and complementary/alternative medicine. *Best Pract Res Clin Rheumatol* 2004;18(4):539-56.
7. Holdcraft LC, Assefi N, Buchwald D. Complementary and alternative medicine in fibromyalgia and related syndromes. *Best Pract Res Clin Rheumatol* 2003;17(4):667-83.
8. Cymet TC. A practical approach to fibromyalgia. *J Natl Med Assoc* 2003;95(4):278-85.
9. Sim J, Adams N. Systematic review of randomized controlled trials of nonpharmacological interventions for fibromyalgia. *Clin J Pain* 2002;18(5):324-36.
10. Targino RA, Imamura M, Kaziyama HH, Souza LP, Hsing WT, Imamura ST. Pain treatment with acupuncture for patients with fibromyalgia. *Curr Pain Headache Rep* 2002;6(5):379-83.
11. Crofford LJ, Appleton BE. Complementary and alternative therapies for fibromyalgia. *Curr Rheumatol Rep* 2001;3(2):147-56.
12. Friedberg F, Sohl S, Schmeizer B. Publication trends in chronic fatigue syndrome: Comparisons with fibromyalgia and fatigue: 1995-2004. *J Psychosom Res* 2007;63(2):143-6.
13. Staud R. Future perspectives: pathogenesis of chronic muscle pain. *Best Pract Res Clin Rheumatol* 2007;21(3):581-96.
14. Nielsen LA, Henriksson KG. Pathophysiological mechanisms in chronic musculoskeletal pain (fibromyalgia): the role of central and peripheral sensitization and pain disinhibition. *Best Pract Res Clin Rheumatol* 2007;21(3):465-80.
15. Mease P, Arnold LM, Bennett R, Boonen A, Buskila D, Carville S, Chappell A, Choy E, Clauw D, Dadabhoy D, Gendreau M, Goldenberg D, Littlejohn G, Martin S, Perera P, Russell IJ, Simon L, Spaeth M, Williams D, Crofford L. Fibromyalgia syndrome. *J Rheumatol* 2007;34(6):1415-25.
16. Abeles AM, Pillinger MH, Solitar BM, Abeles M. Narrative review: the pathophysiology of fibromyalgia. *Ann Intern Med* 2007;146(10):726-34.
17. Tanriverdi F, Karaca Z, Unluhizarci K, Kelestimir F. The hypothalamo-pituitary-adrenal axis in chronic fatigue syndrome and fibromyalgia syndrome. *Stress* 2007;10(1):13-25.
18. Bennett R. Fibromyalgia: Present to future. *Curr Pain Headache Rep* 2004;8(5):379-84.
19. Clauw DJ, Crofford LJ. Chronic widespread pain and fibromyalgia: what we know, and what we need to know. *Best Pract Res Clin Rheumatol* 2003;17(84):685-701.

20. Sarchielli P, Mancini ML, Floridi A, Coppola F, Rossi C, Nardi K, Acciarresi M, Alberto Pini L, Calabresi P. Increased levels of neurotrophins are not specific for chronic migraine: Evidence from primary fibromyalgia syndrome. *J Pain* 2007. Epub Jul 3.
21. Bendtsen L, Norregaard J, Jensen R, Olesen J. Evidence of qualitatively altered nociception in patients with fibromyalgia. *Arthritis Rheum* 1997;40(1):98-102.
22. Henriksson KG. Fibromyalgia – from syndrome to disease. Overview of pathogenetic mechanisms. *J Rehabil Med* 2003;(41 Suppl):89-94.
23. Cook DB, Stegner AJ, McLoughlin MJ. Imaging pain of fibromyalgia. *Curr Pain Headache Rep* 2007;11(3):190-200.
24. Yunus MB. Role of central sensitization in symptoms beyond muscle pain, and the evaluation of a patient with widespread pain. *Best Pract Res Clin Rheumatol* 2007; 21(3):481-97.
25. Nielsen LA, Henriksson KG. Pathophysiological mechanisms in chronic musculoskeletal pain (fibromyalgia): the role of central and peripheral sensitization and pain disinhibition. *Best Pract Res Clin Rheumatol* 2007; 21(3):465-80.
26. Staud R, Vierck CJ, Cannon RL, Mauderli AP, Price DD. Abnormal sensitization and temporal summation of second pain (wind-up) in patients with fibromyalgia syndrome. *Pain* 2001;91(1-2):165-75.
27. Woolf CJ, Salter MW. Plasticity and pain; role of the dorsal horn. In McMahon SB, Koltzenburg M, editors. *Wall and Melzack's Textbook of pain*. London: Elsevier Churchill Livingstone; 2006. p. 91-105.
28. Mense S. The pathogenesis of muscle pain. *Curr Pain Headache Rep* 2003;7(6):419-25.
29. Li J, Simone DA, Larson AA. Windup leads to characteristics of central sensitization. *Pain* 1999;79(1): 75-82.
30. Davies SN, Lodge D. Evidence for involvement of N-methylaspartate receptors in 'wind-up' of class 2 neurones in the dorsal horn of the rat. *Brain Res* 1987;424(2):402-6.
31. Dickenson AH, Sullivan AF. Evidence for a role of the NMDA receptor in the frequency dependent potentiation of deep rat dorsal horn nociceptive neurones following C fibre stimulation. *Neuropharmacology* 1987;26(8):1235-8.
32. Yoshimura Y, Ohmura T, Komatsu Y. Two forms of synaptic plasticity with distinct dependence on age, experience, and NMDA receptor subtype in rat visual cortex. *J Neurosci* 2003;23(16):6557-66.
33. Shum FW, Wu LJ, Zhao MG, Toyoda H, Xu H, Ren M, Pinaud R, Ko SW, Lee YS, Kaang BK, Zhuo M. Alteration of cingulate long-term plasticity and behavioral sensitization to inflammation by environmental enrichment. *Learn Mem* 2007;14(4):304-12.
34. Goldenberg DL. Pharmacological treatment of fibromyalgia and other chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol* 2007;21(3):499-511.
35. Schaible HG, Schmelz M, Tegeder I. Pathophysiology and treatment of pain in joint disease. *Adv Drug Deliv Rev* 2006;58(2):323-42.
36. De Kock MF, Lavand'homme PM. The clinical role of NMDA receptor antagonists for the treatment of postoperative pain. *Best Pract Res Clin Anaesthesiol* 2007;21(1):85-98.
37. Wood PB. A reconsideration of the relevance of systemic low-dose ketamine to the pathophysiology of fibromyalgia. *J Pain* 2006;7(9):611-4.
38. Pozzi A, Muir WW, Traverso F. Prevention of central sensitization and pain by N-methyl-D-aspartate receptor antagonists. *J Am Vet Med Assoc* 2006;228(1):53-60.
39. Gogas KR. Glutamate-based therapeutic approaches: NR2B receptor antagonists. *Curr Opin Pharmacol* 2006;6(1): 68-74.
40. Inturrisi CE. The role of N-methyl-D-aspartate (NMDA) receptors in pain and morphine tolerance. *Minerva Anesthesiol* 2005;71(7-8):401-3.
41. Dickenson AH. A cure for wind up: NMDA receptor antagonists as potential analgesics. *Trends Pharmacol Sci* 1990;11(8):307-9.
42. Quartaroli M, Carignani C, Dal Forno G, Mugnaini M, Ugolini A, Arban R, Bettelini L, Maraia G, Belardetti F, Reggiani A, Trist DG, Ratti E, Di Fabio R, Corsi M. Potent antihyperalgesic activity without tolerance produced by glycine site antagonist of N-methyl-D-aspartate receptor GV196771A. *J Pharmacol Exp Ther* 1999;290(1):158-69.
43. Graven-Nielsen T, Aspegren Kendall S, Henriksson KG, Bengtsson M, Sørensen J, Johnson A, Gerdle B, Arendt-Nielsen L. Ketamine reduces muscle pain, temporal summation, and referred pain in fibromyalgia patients. *Pain* 2000;85(3):483-91.
44. Staud R, Vierck CJ, Robinson ME, Price DD. Effects of the N-methyl D-aspartate receptor antagonist dextromethorphan on temporal summation of pain are similar in fibromyalgia patients and normal control subjects. *J Pain* 2005;6(5):323-32.
45. Klein T, Magerl W, Hanschmann A, Althaus M, Treede RD. Antihyperalgesic and analgesic properties of the N-methyl-d-aspartate (NMDA) receptor antagonist neramexane in a human surrogate model of neurogenic hyperalgesia. *Eur J Pain* 2007. Epub Apr 19.
46. Wang L, Zhang Y, Dai J, Yang J, Gang S. Electroacupuncture (EA) modulates the expression of NMDA receptors in primary sensory neurons in relation to hyperalgesia in rats. *Brain Res* 2006;1120(1):46-53.
47. Sun N, Zou X, Shi J, Liu X, Li L, Zhao L. Electroacupuncture regulates NMDA receptor NR1 subunit expression via PI3-K pathway in a rat model of cerebral ischemia-reperfusion. *Brain Res* 2005;1064(1-2):98-107.
48. Watkins LR, Milligan ED & Maier SF. Glial activation: a driving force for pathological pain. *Trends Neurosci* 2001;24(8):450-5.
49. Watkins LR, Milligan ED, Maier SF. Spinal cord glia: new players in pain. *Pain* 2001;93(3):201-5.
50. Watkins LR, Milligan ED, Maier SF. Glial activation: a driving force for pathological pain. *Trends Neurosci* 2001;24(8):450-5.
51. Gabuzda D, Wang J. Chemokine receptors and virus entry in the central nervous system. *J Neurovirol* 1999;5(6): 643-58.
52. Rock RB, Gekker G, Hu S, Sheng WS, Cheeran M, Lokensgard JR, Peterson PK. Role of microglia in central nervous system infections. *Clin Microbiol Rev* 2004; 17(4):942-64.
53. Watkins LR, Maier SF. The pain of being sick: implications of immune-to-brain communication for understanding pain. *Annu Rev Psychol* 2000;51:29-57.

54. Watkins LR, Maier SF. Immune regulation of central nervous system functions: from sickness responses to pathological pain. *J Intern Med* 2005;257(2):139-55.
55. Watkins LR, Martin D, Ulrich P, Tracey KJ, Maier SF. Evidence for the involvement of spinal cord glia in subcutaneous formalin induced hyperalgesia in the rat. *Pain* 1997;71(3):225-35.
56. Wallace DJ, Linker-Israeli M, Hallegua D, Silverman S, Silver D, Weisman MH. Cytokines play an aetiopathogenic role in fibromyalgia: a hypothesis and pilot study. *Rheumatology (Oxford)* 2001;40(7):743-9.
57. Samad TA, Moore KA, Sapirstein A, Billet S, Allchorne A, Poole S, Bonventre JV, Woolf CJ. Interleukin-1 β -mediated induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature* 2001;410(6827):471-5.
58. Ek M, Engblom D, Saha S, Blomqvist A, Jakobsson PJ, Ericsson-Dahlstrand A. Inflammatory response: pathway across the blood-brain barrier. *Nature* 2001;410(6827):430-1.
59. Raghavendra V, Tanga F, DeLeo JA. Inhibition of microglial activation attenuates the development but not existing hypersensitivity in a rat model of neuropathy. *J Pharmacol Exp Ther* 2003;306(2):624-30.
60. Kang JM, Park HJ, Choi YG, Choe IH, Park JH, Kim YS, Lim S. Acupuncture inhibits microglial activation and inflammatory events in the MPTP-induced mouse model. *Brain Res* 2007;1131(1):211-9.
61. Wang TT, Yuan WL, Ke Q, Song XB, Zhou X, Kang Y, Zhang HT, Lin Y, Hu YL, Feng ZT, Wu LL, Zhou XF. Effects of electro-acupuncture on the expression of c-jun and c-fos in spared dorsal root ganglion and associated spinal laminae following removal of adjacent dorsal root ganglia in cats. *Neuroscience* 2006;140(4):1169-76.
62. Lundeberg T, Lund I, Dahlin L, Borg E, Gustafsson C, Sandin L, Rosen A, Kowalski J, Eriksson SV. Reliability and responsiveness of three different pain assessments. *J Rehabil Med* 2001;33(6):279-83.
63. Stener-Victorin E, Kowalski J, Lundeberg T. A new highly reliable instrument for the assessment of pre- and postoperative gynecological pain. *Anesth Analg* 2002;95(1):151-7.
64. Lund I, Lundeberg T, Kowalski J, Sandberg L, Budh CN, Svensson E. Evaluation of variations in sensory and pain threshold assessments by electrocutaneous stimulation. *Physiother Theory Pract* 2005;21(2):81-92.
65. Lund I, Lundeberg T. Aspects of pain, its assessment and evaluation from an acupuncture perspective. *Acupunct Med* 2006;24(3):109-17.
66. Lund I, Lundeberg T, Kowalski J, Svensson E. Gender differences in electrical pain threshold responses to transcutaneous electrical nerve stimulation (TENS). *Neurosci Lett* 2005;375(2):75-80.
67. Lund I, Lundeberg T. *Do women respond better to acupuncture than men?* Oral presentation at International Symposium of Medical Acupuncture and Related Techniques- 3rd International Medical Congress on Acupuncture (ICMART), Barcelona 1-3 June, 2007
68. Sandberg M, Larsson B, Lindberg LG, Gerdle B. Different patterns of blood flow response in the trapezius muscle following needle stimulation (acupuncture) between healthy subjects and patients with fibromyalgia and work-related trapezius myalgia. *Eur J Pain* 2005;9(5):497-510.
69. Sandberg M, Lindberg LG, Gerdle B. Peripheral effects of needle stimulation (acupuncture) on skin and muscle blood flow in fibromyalgia. *Eur J Pain* 2004;8(2):163-71.
70. Arendt-Nielsen L, Svensson P. Referred muscle pain. Basic and clinical findings. *The Clin J Pain* 2001;17(1):11-9.
71. Hoheisel U, Mense S, Simons DG, Yu X-M. Appearance of new receptive fields in rat dorsal horn neurons following noxious stimulation of skeletal muscle. A model for referral of muscle pain? *Neurosci Lett* 1993;153(1):9-12.
72. Kellgren JH. Observation on referred pain arising from muscle. *Clinical Science (London, England: 1979)* 1938;3:175-190.
73. Inman VT, Saunders JBCM. Referred pain from skeletal structures. *J Nerv Mental Dis* 1944;99:660-667.
74. Simons DG. Muscular Pain Syndromes. In: Friction JR, Awad EA, editors. *Myofascial Pain and Fibromyalgia. Advances in Pain Research and Therapy*. Volume 17. New York: Raven Press; 1990. p. 1-41.
75. Mense S. Nociception from skeletal muscle in relation to clinical muscle pain. *Pain* 1993;54(3):241-289.
76. Mense S. Referral of muscle pain. New aspects. *ASP Journal* 1994;3:1-9.
77. Andersen OK, Spaich EG, Madeleine P, Arendt-Nielsen L. Gradual enlargement of human withdrawal reflex receptive fields following repetitive painful stimulation. *Brain Res* 2005;1042(2):194-204.
78. Kawamata M, Koshizaki M, Shimada SG, Narimatsu E, Kozuka Y, Takahashi T, Namiki A, Collins JG. Changes in response properties and receptive fields of spinal dorsal horn neurons in rats after surgical incision in hairy skin. *Anesthesiology* 2005;102(1):141-51.
79. Andersen OK, Finnerup NB, Spaich EG, Jensen TS, Arendt-Nielsen L. Expansion of nociceptive withdrawal reflex receptive fields in spinal cord injured humans. *Clin Neurophysiol* 2004;115(12):2798-810.
80. Tamburin S, Fiaschi A, Andreoli A, Marani S, Zanette G. Sensorimotor integration to cutaneous afferents in humans: the effect of the size of the receptive field. *Exp Brain Res* 2005;167(3):362-9.
81. Macefield VG. Physiological characteristics of low-threshold mechanoreceptors in joints, muscle and skin in human subjects. *Clin Exp Pharmacol Physiol* 2005;32(1-2):135-44.
82. Kosek E, Ekholm J, Hansson P. Modulation of pressure pain thresholds during and following isometric contraction in patients with fibromyalgia and in healthy controls. *Pain* 1996;64(3):415-23.
83. Staud R, Robinson ME, Price DD. Isometric exercise has opposite effects on central pain mechanisms in fibromyalgia patients compared to normal controls. *Pain* 2005;118(1-2):176-84.
84. Suzuki R, Rygh LJ, Dickenson AH. Bad news from the brain. Descending 5-HT pathways that control spinal pain processing. *Trends Pharmacol Sci* 2004;25(12):613-7.
85. Vierck CJ Jr. Mechanisms underlying development of spatially distributed chronic pain (fibromyalgia). *Pain* 2006;124(3):242-63.
86. Ren K, Dubner R. Descending modulation in persistent pain: an update. *Pain* 2002;100(1-2):1-6.
87. Suzuki R, Rahman W, Hunt SP, Dickenson AH. Descending facilitatory control of mechanically evoked responses is

- enhanced in deep dorsal horn neurones following peripheral nerve injury. *Brain Res* 2004;1019(1-2):68-76.
88. Jones AKP. The role of the cerebral cortex in pain perception. In: Justins DM, editor. *Refresher Course Syllabus. Pain 2005 – An updated review*. Seattle: IASP Press; 2005. p. 59-68.
 89. Deluze C, Bosia L, Zirbs A, Chantraine A, Vischer TL. Electroacupuncture in fibromyalgia: results of a controlled trial. *BMJ* 1992;305(6864):1249-52.
 90. Thomas M, Lundberg T. Importance of modes of acupuncture in the treatment of chronic nociceptive low back pain. *Acta Anaesthesiol Scand* 1994;38(1):63-9.
 91. Thomas M, Lundberg T, Bjork G, Lundstrom-Lindstedt V. Pain and discomfort in primary dysmenorrhea is reduced by preemptive acupuncture or low frequency TENS. *Eur J Phys Med Rehabil* 1995;5(3):71-6.
 92. Dieter HH, Muckter H. Assessment of so called organic trace compounds in drinking water from the regulatory, health and aesthetic-quality points of view, with special consideration given to pharmaceuticals. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2007;50(3):322-31.
 93. Ritter L, Solomon K, Sibley P, Hall K, Keen P, Mattu G, Linton B. Sources, pathways, and relative risks of contaminants in surface water and groundwater: a perspective prepared for the Walkerton inquiry. *J Toxicol Environ Health A* 2002;65(1):1-142.
 94. Stackelberg PE, Furlong ET, Meyer MT, Zaugg SD, Henderson AK, Reissman DB. Persistence of pharmaceutical compounds and other organic wastewater contaminants in a conventional drinking-water-treatment plant. *Sci Total Environ* 2004;329(1-3):99-113.
 95. Liebig M, Moltmann JF, Knacker T. Evaluation of measured and predicted environmental concentrations of selected human pharmaceuticals and personal care products. *Environ Sci Pollut Res Int* 2006;13(2):110-9.
 96. Krishnan K, Paterson J, Williams DT. Health risk assessment of drinking water contaminants in Canada: the applicability of mixture risk assessment methods. *Regul Toxicol Pharmacol* 1997;26(2):179-87.
 97. Wollin K, Dieter HH. Toxicological guidelines for monocyclic nitro-, amino- and aminonitroaromatics, nitramines, and nitrate esters in drinking water. *Arch Environ Contam Toxicol* 2005;49(1):18-26.
 98. Bartel H, Kruger W, Mendel B, Suhr R. The drinking water ordinance—successful or requiring revision? *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2007;50(3):265-75.
 99. Webb S, Ternes T, Gibert M, Olejniczak K. Indirect human exposure to pharmaceuticals via drinking water. *Toxicol Lett* 2003;142(3):157-67.
 100. Falconer IR. Are endocrine disrupting compounds a health risk in drinking water? *Int J Environ Res Public Health* 2006;3(2):180-4.
 101. Zuccato E, Castiglioni S, Fanelli R, Reitano G, Bagnati R, Chiabrando C, Pomati F, Rossetti C, Calamari D. Pharmaceuticals in the environment in Italy: causes, occurrence, effects and control. *Environ Sci Pollut Res Int* 2006;13(1):15-21.
 102. Radel U, Puchert W, Suchenwirth R. Surveillance of drinking water supply systems on markets and in vehicles. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2007;50(3):332-9.
 103. Jones OA, Lester JN, Voulvoulis N. Pharmaceuticals: a threat to drinking water? *Trends Biotechnol* 2005;23(4):163-7.